

$$[1 - C/A] \cos^2 \theta_1 = [1 - C/B] \cos^2 \theta_0 \quad (1)$$

The increase in θ from θ_0 to θ_1 as the twist angle α increases from 0 to 90° will be greatest when the arms are abducted (wide arms). Let the values A_0, B_0, C_0, θ_0 and α_0 correspond to this body configuration. The decrease in θ from θ_0 to θ_1 as α increases from 90° to 180° will be smallest when the arms are adducted (touching the body). Let the values A_1, B_1, C_1, θ_1 and α_1 correspond to this configuration. Suppose that during a half twist the arms are rapidly adducted at the quarter twist position so that $\theta_1 = \theta_0$ and then rapidly abducted again at the half twist position. Two applications of equation (1) show that the tilt angle increases from θ_0 to θ_1 such that:

$$\cos \theta_1 = k \cdot \cos \theta_0$$

where:

$$k^2 = \frac{[1 - C_1/A_1][1 - C_0/B_0]}{[1 - C_1/B_1][1 - C_0/A_0]}$$

(11/1/2023)

(11/1/2023)






Figure 5: Simulation of a double-full-out straight.

to the full-out straight and double-full-out straight. These movements are used in dismounts from the gymnastics apparatus and in trampoline routines.

References

- Yeadon, M. (1986). The biomechanics of twisting somersaults. In *Proceedings of the North American Conference on Biomechanics*, pages 33–34.
- Yeadon, M. R. (1993a). The biomechanics of twisting somersaults. Part I: Rigid body motions. *Journal of Sports Sciences*, 11:187–198.
- Yeadon, M. R. (1993b). The biomechanics of twisting somersaults. Part II: Contact twist. *Journal of Sports Sciences*, 11:199–208.